arXiv:math/0505297v2 [math.GT] 17 May 2005

On transverse Hopf links

Ichiro Torisu

Naruto University of Education, 748, Nakajima, Takashima, Naruto-cho, Naruto-shi, 772-8502 Japan
e-mail: torisu@naruto-u.ac.jp

Date: May 13, 2005.

Abstract

We classify transverse Hopf links in the standard contact 3-space up to transverse isotopy in terms of their components' self-linking number.

1 The statement of result

A *contact structure* on a 3-manifold is a completely non-integrable 2-plane field on it. Let ξ_0 be the standard contact structure on 3-space $R^3 = \{(x, y, z)\}$, that is, a 2-plane field on R^3 defined by the kernel of the 1-form $dz - ydx$. A *transverse knot/link* is an oriented knot/link which is everywhere transverse to the contact structure. Throughout, we always assume a transverse knot/link is positively oriented to ξ_0.

Let T be a transverse knot in the standard contact 3-space (R^3, ξ_0) and v be a non-singular vector field in ξ_0 on R^3. Then the *self-linking number* (aka. *Bennequin number*) $sl(T)$ of T is defined to be the linking number of T and a knot slightly pushed off T along v. Note that $sl(T)$ is independent of choices of v and an orientation of T, $sl(T)$ is invariant under transverse isotopy of T and that $sl(T)$ equals the writhe in a generic projection of T to xz-plane. In [1], D. Bennequin showed that $sl(T)$ is bounded by the negative Euler characteristic of a Seifert surface for T.

In [3], Y. Eliashberg classified topologically trivial transverse knots in (R^3, ξ_0) up to transverse isotopy in terms of their self-linking number. Therefore transverse knots featured in Figure 1 give a complete list of transverse unknots in (R^3, ξ_0), where the diagrams are drawn on xz-plane and the self-linking number is a negative odd integer.

In [2], J. Birman–M. Wrinkle classified topologically trivial transverse *links* in (R^3, ξ_0) up to transverse isotopy in terms of their components' self-linking number.



Figure 1: A transverse unknot

Recently, J. Etnyre–K. Honda classified transverse torus knots and transverse figure-eight knots in terms of their self-linking number ([4], [6]). But there is not much result of transverse links with more than one component.

In this paper, we will classify transverse *Hopf links* in the standard contact 3-space. For the proof, we will apply the Eliashberg's original arguments of analyzing a characteristic foliation of a disk to an immersed disk spanned by a Hopf link. Remark that in [8], the author also used the arguments for study of transverse *doubled knots.*

Let $H_+ = l_1^+ \cup l_2^+$ and $H_- = l_1^- \cup l_2^-$ be the positive and negative Hopf links in R^3, where l_i^+ and l_i^- are the components and the linking numbers $lk(l_1^+, l_2^+)$ and $lk(l_1^-, l_2^-)$ are -1 and 1, respectively.

Theorem 1. Two transverse Hopf links in (R^3, ξ_0) are transversely isotopic to each other if and only if their topological type H_+/H_- and the pair of the self-linking number of their components agree.

Remark 1. (i) By Theorem 1, transverse links featured in Figure 2 give a complete list of transverse Hopf links in (R^3, ξ_0) up to transverse isotopy, where the diagrams are drawn on xz-plane.
(ii) The statement of Theorem 1 implies that for a transverse Hopf link in (R^3, ξ_0) with the same self-linking number of two component, there is a transverse isotopy which exchanges the components.
(iii) After writing this paper, the author was pointed that the part for positive Hopf links is proved in the Etnyre's positive torus knots paper [4] though he doesn't state that it was proved.



Figure 2: transverse Hopf links



Figure 3: A clasp disk

2 Characteristic foliations on immersed disks

Let $H = l_1 \cup l_2$ be an oriented Hopf link in R^3, where l_i is the component. Let $\tilde{D}_1 \amalg \tilde{D}_2$ be a disjoint union of two 2-disks. Then there exists an immersion $f : \tilde{D}_1 \amalg \tilde{D}_2 \to R^3$ such that (i) $f_{|\partial\tilde{D}_1 \amalg \partial\tilde{D}_2} : \partial\tilde{D}_1 \amalg \partial\tilde{D}_2 \to H$ is a diffeomorphism and $f(\partial\tilde{D}_i) = l_i$ (ii) $f^{-1}(\Omega)$ consists of a disjoint union of two arcs $\tilde{\sigma}_1 \amalg \tilde{\sigma}_2$ such that each $\tilde{\sigma}_i$ joins a point of $\partial\tilde{D}_i$ to a point in $Int\tilde{D}_i$, where $\Omega = \{x \in f(\tilde{D}_1 \amalg \tilde{D}_2) | |f^{-1}(x)| \geq 2\}$. Put $D = f(\tilde{D}_1 \amalg \tilde{D}_2)$, $D^1 = f(\tilde{D}_1)$ and $D^2 = f(\tilde{D}_2)$. Then $D = D^1 \cup D^2$ is called a *clasp disk* for H and $\sigma = D^1 \cap D^2 = f(\tilde{\sigma}_1) = f(\tilde{\sigma}_2)$ is called the *clasp singularity* (Figure 3). Throughout, we use the notation ∂D as $f(\partial\tilde{D}_1 \amalg \partial\tilde{D}_2)$ (ie., $\partial D = H$). Then D is an oriented immersed disk in R^3 such that the induced orientation of ∂D agrees with the orientation of H.

Further, suppose H is a transverse link in (R^3, ξ_0). Let D_{ξ_0} be the *characteristic foliation* of D, that is, the singular foliation defined by $TD \cap \xi_0$ on D. Remark that D_{ξ_0} is orientable. Throughout, we orient D_{ξ_0} in such a way that trajectories of D_{ξ_0} exit through ∂D.

Generically D_{ξ_0} satisfies the following:

- D is tangent to ξ_0 at a finite set Σ in $D - \partial D$ and all singular point in Σ is an



Figure 4: A characteristic foliation

elliptic point or a *hyperbolic* point according to the index (1 or -1, respectively),

- there is no separatrix connection between hyperbolic points.

See Figure 4. In this paper, we call a clasp disk satisfying these conditions a *generic* clasp disk.

And we can distinguish between *positive* and *negative* points of Σ depending on whether the orientations of D and ξ_0 coincide at these points or not.

Denote the restriction of D_{ξ_0} to D^i by $D^i_{\xi_0}$ $(i = 1, 2)$. That is, $D^i_{\xi_0}$ is the characteristic foliation of D^i. Let e^i_+ (e^i_-, resp.) and h^i_+ (h^i_-, resp.) be the number of positive (negative, resp.) elliptic and positive (negative, resp.) hyperbolic points in $D^i_{\xi_0}$, respectively. Then from $D^i_{\xi_0}$, we can compute the self-linking number of each component l_i of ∂D as follows.

Proposition 1 ([1]). $sl(l_i)$ is equal to $-e^i_+ + e^i_- + h^i_+ - h^i_-$ $(i = 1, 2)$.

A curve in (R^3, ξ_0) is called *Legendrian* if it is everywhere tangent to ξ_0.

Proposition 2. For any transverse Hopf link H in (R^3, ξ_0), there is a generic clasp disk D for H such that the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ. In particular, there is no singular point of the characteristic foliation D_{ξ_0} along σ.

Proof of Proposition 2. Take a *Legendrian push off* L of H such that L is C^0-close to H and each component of L is a closed leaf on the boundary of a standard neighbourhood of each component of H ([3], [5]). Note that a Legendrian isotopy of L induces a transverse isotopy of H ([3], [5]). Without loss of generality, we assume L is given by a *front projection* on xz-plane in R^3. Then after Legendrian Reidemeister moves to L , if necessary, we can find a Legendrian clasp part between two component



Figure 5: Legendrian clasps



Figure 6:

of L as in Figure 5. Then it is easy to see that there is a clasp disk D_L for L such that along the interior of its clasp singularity σ_L, there is no tangency of D_L to ξ_0, where σ_L is indicated by the dotted segment in Figure 5. From D_L, we can easily make the desired clasp disk for H. This completes the proof.

For the remainder of this section, we prove the following.

Theorem 2. Let H be a transverse Hopf link in (R^3, ξ_0). Then there exists a generic clasp disk $D = D^1 \cup D^2$ for H satisfying the following (Figure 6):

- the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ,
- $e^i_- = h^i_+ = 0$ in $D^i_{\xi_0}$ $(i = 1, 2)$,
- σ is included in an unstable trajectory from a positive elliptic point in $D^i_{\xi_0}$ $(i = 1, 2)$.

Proof of Theorem 2. By Proposition 2, there is a generic clasp disk D for H such that the clasp singularity σ of D is Legendrian and ξ_0 is transverse to D along σ.

Then the next lemma due to E. Giroux is a basic tool of a modification of D_{ξ_0} which allows us to cancel or create the pairs of an elliptic point and a hyperbolic point of the same sign in D_{ξ_0}.

Lemma 1 (Elimination lemma [3], [7]). Let $p, q \in D - \sigma$ be elliptic and hyperbolic points of D_{ξ_0} of the same sign. Let γ be a trajectory of D_{ξ_0} which joins p and q and is disjoint from σ. Then there exists a C^0-small isotopy $f_t : \tilde{D_1} \amalg \tilde{D_2} \to R^3, t \in [0,1]$, which starts with $f_0 = f$, is fixed on γ and outside of a neighbourhood $U \supset \gamma$ and such that the characteristic foliation of $f_1(\tilde{D_1} \amalg \tilde{D_2})$ has no singularity in $U \cap f_1(\tilde{D_1} \amalg \tilde{D_2})$.

We first eliminate all negative elliptic point in D_{ξ_0}. Let p be a negative elliptic point in D_{ξ_0}. If there is a negative hyperbolic point connected with p by a trajectory in D_{ξ_0}, then we cancel the pair by Lemma 1. If otherwise, as in the proof of Theorem 4.1.1 in [3], after creating the pairs of positive elliptic and positive hyperbolic points, if necessary, we have a limit cycle in D_{ξ_0} which is disjoint from both ∂D and σ. This contradicts the Bennequin inequality.

Next, we eliminate all positive hyperbolic point in D_{ξ_0}. Let q be a positive hyperbolic point in D_{ξ_0}. Then by our assumptions, a stable separatrix of q necessarily connects q with a positive elliptic point. Again we cancel the pair by Lemma 1.

Finally, by the genericity of D_{ξ_0}, we can modify D so that σ is not included in an unstable trajectory from a negative hyperbolic point. This completes the proof of Theroem 2.

3 Proof of Theorem 1

Recall from [4] that a transverse link K is a *single stabilization* of a transverse link K' if $K = \beta \cup A, K' = \beta \cup A'$ and $A \cup A'$ cobounds a disk such that (i) the characteristic foliation of the disk admits only one positive elliptic and one negative hyperbolic points as singularities (ii) the interior of the disk is disjoint from K' (iii) $sl(K) = sl(K') - 2$ (Figure 7).



Figure 7:

For the proof of Theorem 1, we need the following Etnyre's theorem in [4].

Theorem 3 ([4]). Let (K, l), (K', l'), (C, d) and (C', d') be the pairs of transverse links and their one components. Suppose transverse links K and K' are single stabilizations of transverse links C and C' and the stabilizations are done on the pairs (l, d) and (l', d'), respectively. Then (K, l) is transversely isotopic to (K', l') pairwisely if (C, d) is transversely isotopic to (C', d') pairwisely.

In [4], Etnyre originally gave a proof for the case where K, K', C and C' are *knots*, but the proof is directly applicable to our links case.

By Theorem 2 and Theorem 3, we only need to prove that transverse links in the link type H_+/H_- such that the self-linking number of each component is maximal (i.e. -1) are unique up to transverse isotopy. In particular, for such a transverse Hopf link, there is a transverse isotopy which exchanges the components.

The next lemma is easy to prove by considering front projections.

Lemma 2. Suppose c and c' are oriented Legendrian embedded arcs in (R^3, ξ_0). Then there is a contact isotopy of (R^3, ξ_0) which sends c to c' with their orientations.

Let H and H' be such transverse Hopf links. Then, there exist generic clasp disks D snd D' for H and H' satisfying the statement of Theorem 2, respectively. Since the self-linking number of each component of H and H' is maximal, the characteristic foliations D_{ξ_0} and D'_{ξ_0} admit exactly two positive elliptic points as singularities, respectively. Let c and c' be Legendrian arcs embedded in D_{ξ_0} and D'_{ξ_0} such that the boundaries of c and c' equal the two positive elliptic points of D_{ξ_0} and D'_{ξ_0} and that c and c' contain the clasp singularities of D and D', respectively. Then, for any choice of orientations of c, c', by Lemma 2 there is a contact isotopy of (R^3, ξ_0) which sends c to c' with the chosen orientations. Furthermore, since D_{ξ_0} is orientation preservings diffeomorphic to D'_{ξ_0}, by using D and D', we can easily make a desired transverse isotopy from H to H'. This completes the proof of Theorem 1.

References

[1] D. Bennequin, *Entrelacements et équations de Pfaff*, Astérisque, **107-108** (1983) 87–161.

[2] J. Birman–N. Wrinkle, *On transversally simple knots*, J. Differential Geometry, **66** (2000) 325–354.

[3] Y. Eliashberg, *Legendrian and transversal knots in tight contact 3-manifolds*, Topological Methods in Modern Mathematics, (1993) 171–193.

[4] J. Etnyre, *Transversal torus knots*, Geomtery and Topology, **3** (1999) 253–268.

[5] J. Etnyre, *Legendrian and transversal knots*, to appear in the Handbook of Knot Theory (available at http://front.math.ucdavis.edu/math.SG/0306256).

[6] J. Etnyre–K. Honda, *Knots and Contact geometry I: torus knots and the figure eight*, J. Symplectic Geometry, **1** (2001) 63–120.

[7] E. Giroux, *Convexité en topologie de contact*, Comment. Math. Helvetici, **66** (1991) 637–677.

[8] I. Torisu, *Transverse doubled knots and clasp disks*, preprint (available at http://front.math.ucdavis.edu/math.GT/0505023).